

# Lecitop



## DESCRIPTION

**LECITOP** is a natural product based on soya lecithin, which contains substances such as phospholipids, essential fatty acids, choline, inositol, phosphorus and vitamins.

Lecithins are classified as Basic Substances with FUNGICIDE action, according to SANCO report 12798/2014 and are regulated in Article 23 of Regulation (EC) 1107/2009, concerning the placing of plant protection products on the market.

## FEATURES


**LECITOP**, hardens the cell wall of the crops, at the same time as it gives plasticity to the fruits, preventing physiopathies and improving consistency. It also exerts a hydrophobic activity on the leaf surface, preventing relative leaf surface, preventing relative humidity from resting on the plant surface and reducing the proliferation of aerial fungi.

**It can be used generically against the vast majority of fungi that cause damage**

- It is an effective antioxidant providing longevity and turgor in fruits and vegetables.
- Improves skin anomalies by giving colour and shine, thus making the fruit more attractive to the eye.
- It heals small physiopathies and wounds on fruits.
- Improves plant health in general, reducing fungal and bacterial attacks.

**Product SUITABLE for use in organic farming in accordance with Regulation (EU) 2018/848**

## PROPERTIES



## DECLARED CONTENT

| BASIC SUBSTANCE USEFUL IN PLANT PROTECTION | w/w                    |
|--------------------------------------------|------------------------|
| Soy Lecithins                              | 30 %                   |
| Density                                    | 0,94 g/cm <sup>3</sup> |
| pH                                         | 7                      |

## OPTIONAL INFORMATION

BASIC SUBSTANCE USEFUL IN PLANT PROTECTION IN ACCORDANCE WITH REGULATION (EC) No 1107/2009

# Lecitop

## RECOMMENDED APPLICATION PER CROP

| CROP                                                | Disease                                                      | Dose        | Application                                                                                                           |
|-----------------------------------------------------|--------------------------------------------------------------|-------------|-----------------------------------------------------------------------------------------------------------------------|
| <b>Apple tree</b>                                   | Mildiu "Podosphaera leucotricha"                             | 750 cc/HI   | <b>3 to 12 applications</b><br>Bud swelling to 90% fruit every 5 days                                                 |
| <b>Peach tree</b>                                   | Taphrina deformans<br>Lepra ó abolladura                     | 750 cc/HI   | <b>3 to 12 applications</b><br>Bud swelling to 90% fruit<br>fruit to 90% - every 5 days                               |
| <b>Vineyard</b>                                     | Mildiu,<br>"Pasmopara viticola", Oídio<br>"Erysiphe necator" | 750 cc/HI   | <b>3 to 12 applications</b><br>leaf not unfolded until<br>berry softening<br>Every 5 days                             |
| <b>Vegetables</b><br>Cucumber<br>Courgette<br>Melon | Oídio<br>"Podosphaera xanthii Oídio"                         | 1.500 cc/HI | <b>2 to 12 applications</b><br>Cotyledons until full<br>maturity. Interval: 5 days                                    |
| <b>Potato</b>                                       | Mildiu<br>"Phytophthora infestans"                           | 1.800 cc/HI | <b>3 to 12 applications</b><br>9 leaves to maturity.<br>Every 5 days                                                  |
| <b>Lettuce</b>                                      | Oídio<br>"Erysiphe cichoracearum"                            | 1.500 cc/HI | <b>2 applications</b><br>Cotyledons until maturity.<br>Interval: 7 days                                               |
| <b>Tomato</b>                                       | Mildiu<br>"Phytophthora infestans"                           | 1.500 cc/HI | <b>2 to 6 applications</b><br>Cotyledons until maturity.<br>Interval: 5 days                                          |
| <b>Carrot</b>                                       | "Leveillula taurica Mildiu"                                  | 1.800 cc/HI | <b>4 applications</b> from 9 or<br>more leaves unfolded until<br>fruit ripening BBCH 19-<br>BBCH89. Interval: 14 days |
| <b>Hornamental</b>                                  | Against various fungi including<br>including mildew.         | 750 cc/HI   | <b>3-12 applications</b> during<br>the whole crop cycle.<br>Interval: 5 days                                          |



CARING FOR AGRICULTURE  
WE ALSO CARE FOR THE PLANET

DUNYA AGRONOMIC SOLUTIONS S.L. Pinseque (Zaragoza) Spain - Tel/WhatsApp :+34 876 00 95 74- info@dunya.es