



Sanitur

DESCRIPTION

SANITUR, has been developed under RD(CE)1.107/2.009 from "Urtica spp" to provide a natural and low environmental impact tool against diseases such as downy mildew, potato blight or pests such as aphids.

SANITUR is composed of Nettle extract, indicated as an insecticide, fungicide and acaricide product.

FEATURES

SANITUR, is generated by extraction in an aqueous medium, it provides numerous nutrients and some active substances such as flavonoids, tannins, organic acids, etc., which promote better and healthy plant development.

These organic compounds applied on plants complement nutrition, having an active response when applied.

SANITUR, used continuously throughout the plant's life cycle, it can keep pathogenic insect populations away from the crop and reduce their impact on yields.

It is particularly effective on small insects such as aphids and mites.

Product **SUITABLE** for use in organic farming in accordance with Regulation (EU) 2018/848

PROPERTIES



DECLARED CONTENT

BASIC SUBSTANCE USEFUL IN PLANT PROTECTION		Quantity
Urtica spp		15 gr/litro
Density		1,05g/cm ³
pH		5-6

FORM OF APPLICATION

FOLIAR APPLICATION: The average application rate is between 50 to 100 cc/Hl.
For more information see authorised crops, pests and safety periods.

DUN-FM-2021

DUNYA AGRONOMIC SOLUTIONS S.L. Pinseque (Zaragoza) Spain - Tel/WhatsApp :+34 876 00 95 74- info@dunya.es

Sanitur

Crop and Objective	Dose	Mode of Application
AGAINST APHIDS /Apple tree; Pear tree; Plum; Peach; Currant (white and red); Walnut tree red and blackcurrant); Walnut tree; Cherry tree; Lettuce (outdoor); Endive; Chicory; Ornamental trees and shrubs (not for consumption); Green and dried beans dried beans; Potatoes.	10 l/Hl Broth of 300 l Ha	1-5 applications , wetting the leaves and stems (in the presence of aphids). Security period: 7 days
LA CARPOCAPSA Apple tree.	10 l/Hl Broth of 300 l Ha	1-3 applications (interval between applications 7-21 days). Security period: 7 days
ALTERNARIOSIS — Apple; Pear tree peach; cherry; broccoli; broccoli Cauliflower (white and green); Brussels sprouts; Chinese cabbage; RosettePakchoi and Komatsuna; Curly kale; Cabbage; RadishRape; Radish; Gherkins; Courgettes; Melons; Pumpkin.	10 l/Hl Broth of 300 l Ha	1-6 applications during the summer Security period: 7 days
MONOLIOSE — Apple; Pear; Peach; Cherry.	10 l/Hl Broth of 300 l Ha	1-6 applications Security period: 7 days
BOTRYTIS — Apple tree; Plum tree; Peach; Cherry.	10 l/Hl Broth of 300 l Ha	1-6 applications Security period: 7 days
POST-HARVEST DISEASES Apple tree; Plum tree; Peach tree Cherry	10 l/Hl Broth of 300 l Ha	1-6 applications Security period: 7 days
OÍDIUM — Cucumber; Gherkin; Pumpkin and Courgette.	10 l/Hl Broth of 300 l Ha	1-6 applications Security period: 7 days
MIDIU — Grapes and table grapes Potatoes.	10 l/Hl Broth of 300 l Ha	1-6 applications during the summer Security period: 7 days.
ACAROS TETRANYCHIDAE — Grapes and table grapes; Dried and green beans.	10 l/Hl Broth of 300 l Ha	1-6 applications during the summer Security period: 7 days.
ALTICES — Kohlrabi; Kale, Broccoli; Cauliflower; Brussels sprouts; Cabbages; Chinese cabbage; Pakchoi; Rosettepakchoi and komatsuna; Rapeseed and Radish.	10 l/Hl Broth of 300 l Ha	1-6 applications during the summer Security period: 7 days.
DIAMONDBACK MOTH(PDD) Kohlrabi; Kale, Broccoli; Cauliflower; Brussels sprouts; Cabbages; Chinese cabbage; Pakchoi; Rosette pakchoi and komatsuna; Rapeseed and Radish	10 l/Hl Broth of 300 l Ha	1-6 applications during the summer Security period: 7 days.

DUN-FM-2021

DUNYA AGRONOMIC SOLUTIONS S.L. Pinseque (Zaragoza) Spain - Tel/WhatsApp :+34 876 00 95 74- info@dunya.es